

Childhood Immunization Schedule

Approved by the Advisory Committee on Immunization Practices,
American Academy of Pediatrics, and the American Academy of Family Physicians

PacifiCare®

A UnitedHealthcare Company

AGE	Birth	2 mos	4 mos	6 mos	12-15 mos	15-18 mos	2 yrs
Hepatitis B	■	■	■	■	■		
DTaP (diphtheria, tetanus, pertussis)		■	■	■		■	
Hib (haemophilus influenzae type b)		■	■	■	■		
IPV (inactivated polio virus)		■	■	■	■		
PCV ¹ (pneumococcal conjugate vaccine)		■	■	■	■		
Rotavirus (oral vaccine)		■	■	■			
MMR (measles, mumps, rubella)					■		
Varicella (chickenpox)						■	
Influenza ² (flu shot)					■		
Hepatitis A ³						■	■

1. PCV (Prevnar™) helps to protect against streptococcus pneumoniae (PNC) which is the most common bacterial cause of acute otitis media (ear infections), community-acquired pneumonia, and bacteremia (bacteria in the blood).
2. Influenza vaccine (flu shot) is recommended annually (during flu season) for children age 6-23 months. For children 24 months and older, check with your child's doctor for recommendations.
3. Hepatitis A series is recommended at 1-2 years of age, with a booster administered 6-18 months after the 1st dose.